TEPEZZA®
teprotumumab-trbw

TEPEZZA is for adults with Thyroid Eye Disease (TED), no matter if they've had TED for months or years

TEPEZZA as a treatment option for TED

Randi, a real TEPEZZA patient.
Randi was compensated for her time.

Learn about a treatment that has been used by

OVER 25,000 PATIENTS*

*Based on 25,062 individual patients enrolled in Amgen By Your Side who received at least one dose of TEPEZZA from 02/05/2020 to 11/14/2025. TED diagnosis is assumed based on initiation of TEPEZZA.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TEPEZZA?

Infusion reactions can happen during or within 24 hours after your infusion of TEPEZZA. If you have a reaction while receiving TEPEZZA, your doctor or nurse will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction, your doctor may stop your treatment completely.

Please see additional Important Safety Information for TEPEZZA throughout and on pages 12-13.

What is Thyroid Eye Disease (TED)?

TED is an ongoing autoimmune condition that affects the eyes and can get worse over time

TED looks different for everyone:

- Symptoms can appear at any time
- Symptoms can range from mild to severe
- New or previously experienced symptoms may temporarily come back or get worse, which is called a flare
- About 1 out of 6 people with TED experience flares, regardless of how their TED is treated
- Flares can occur years after early symptoms stopped or lessened

The Tiresias Font
Designed for Readability

TED can make reading hard. So we're dedicated to making our materials more accessible. At Amgen, we know vision issues caused by TED, like double vision, can make it hard to read or focus. That's why we tested which fonts were easiest to see. The result is the font you see here, called "**Tiresias**." **It was designed specifically for people with low vision.**

TEPEZZA targets a root cause of TED

TEPEZZA, the first FDA-approved medicine for TED, is a non-surgical treatment designed to target a root cause of TED. TED is an autoimmune condition in which your body's immune system mistakenly attacks the tissue behind your eyes, causing inflammation and swelling. TEPEZZA was designed to block the body's response to this autoimmune condition, reducing inflammation and swelling of muscle and fat tissue.





The exact nature of how TED develops and progresses is not entirely understood, therefore, precisely how TEPEZZA works is not fully established.

IMPORTANT SAFETY INFORMATION (cont'd)

Tell your doctor or nurse right away if you have any of these symptoms during or after your treatment with TEPEZZA:

- High blood pressure
- Fast heartbeat
- Redness of the face/Feeling hot
- Difficulty breathing
- Headache
- Muscle pain

Please see additional Important Safety Information for TEPEZZA throughout and on pages 12-13.

Learn about the most common symptoms of TED

TED can cause symptoms like:

- Eye pain and/or swelling
- Double vision
- Bulging eyes
- Misaligned eyes
- Pulled-back eyelids
- Blurry vision
- Red, dry, gritty eyes
- Watery, teary eyes

These symptoms can lead to impacts on your daily life like:

- Trouble with daily activities, like walking or driving, due to impact on vision
- Feeling that your appearance has changed, or that your eyes look different in photos
- Feeling less confident
- Appearing in photos less often or trying to mask changes in your appearance
- Difficulty completing work-related tasks or activities, such as reading or viewing screens, due to impact on vision

TEPEZZA may be able to help

With TED, it's not just about your eye symptoms—it's about how those symptoms can impact the activities of your daily life. TEPEZZA may help reduce that impact.

TEPEZZA®
teprotumumab-trbw

Even ONE TED SYMPTOM + Even ONE IMPACT on daily life = ASK ABOUT TEPEZZA

A conversation with your doctor may be able to help

Hear from Randi,

a real TEPEZZA patient

"When that double vision came on… looking at my computer screen, driving… those were things I no longer could do. I completed all eight infusions and the symptoms have never returned."

— Randi C.

Randi has been compensated for her time.

Ready to start the conversation with your doctor? Get a Conversation Guide

IMPORTANT SAFETY INFORMATION (cont'd)

TEPEZZA may worsen inflammatory bowel disease (IBD), such as Crohn's disease or ulcerative colitis, even if you don't have a diagnosis of IBD before starting TEPEZZA treatment. Tell your doctor right away if you have worsening IBD symptoms, which may include diarrhea with stomach pain or cramps, blood in your stools, sudden urgency to have a bowel movement, or feeling like you need to have a bowel movement even when there's little or no stool to pass.

Please see additional Important Safety Information for TEPEZZA throughout and on pages 12-13.

TEPEZZA®
teprotumumab-trbw

TEPEZZA has been proven effective in treating TED

In a clinical study of recently diagnosed TED patients*

8 out of 10 people who took TEPEZZA saw **reduced eye bulging** vs 1 out of 10 people who took placebo‡

Individual results may vary.
*24-week clinical trial included 41 patients in the TEPEZZA group who had TED for 9 months or less.
‡Eye bulging was reduced by at least 2 mm.

You could see results after your second infusion‡

- With TEPEZZA, some people began to see less eye bulging in as few as 6 weeks, with improvements continuing for the full course of 8 infusions

‡At Week 6, there was an average of -2 mm of reduced proptosis for patients receiving TEPEZZA compared to an average of -0.38 mm reduction for patients receiving placebo.

Real results with TEPEZZA





Real patients treated with TEPEZZA. Individual results may vary.

See more clearly by improving double vision

In clinical studies of recently diagnosed TED patients§

7 out of 10 people who took TEPEZZA saw **improvements in double vision** vs 3 out of 10 people who took placebo

Individual results may vary.
§Two 24-week clinical trials included 84 patients in the TEPEZZA group who had TED for 9 months or less.

- In those same clinical studies,** more than 5 out of 10 people taking TEPEZZA saw their double vision completely go away, versus an average of 2.5 out of 10 people taking placebo

**Two studies were conducted throughout 24 weeks each and followed patients taking TEPEZZA who were recently diagnosed with moderate-to-severe TED.

IMPORTANT SAFETY INFORMATION (cont'd)

TEPEZZA may cause an increase in your blood sugar. Before starting treatment with TEPEZZA, tell your doctor if you are currently being treated for diabetes, know your blood sugar is high, or have been diagnosed with diabetes. It is important for you to take your treatments and follow an appropriate diet for glucose control as prescribed by your doctor.

TEPEZZA may cause severe hearing problems including hearing loss, which in some cases may be permanent. Tell your doctor if you have any signs or symptoms of hearing problems or changes in hearing.

Please see additional Important Safety Information for TEPEZZA throughout and on pages 12-13.

TEPEZZA®
teprotumumab-trbw

Get multi-symptom relief

In clinical studies, 2 out of 3 people who completed treatment with TEPEZZA had little to no:

Eye pain
including pain behind the eye and when moving the eye

Redness
of eye and eyelid

Swelling
of the eye and eyelid

In combined results from two studies, TED patients' levels of inflammation were recorded using their Clinical Activity Score (CAS), which is a tool that doctors use to assess their patients' TED symptoms. It evaluates for the presence of 7 symptoms in the eyes and surrounding area: pain that happens for no apparent reason; pain due to looking at something for too long; eyelid swelling; eyelid redness; redness of the eye itself; swelling of the eye, and inflammation in the inner corner of the eye. The higher the CAS number (from 0 to 7), the more serious the impact of TED is seen to be.

In patients with a CAS of at least 4 at the start of treatment, by the end of treatment 62% (52 of 84) of those treated with TEPEZZA had a CAS of 0 or 1 (seen as inactive disease state) compared to 22% (18 of 87) of those who were treated with placebo.

Each of the 7 symptoms is counted equally toward a single overall score, even though that may not reflect their impact to the individual patient. For example, if a patient has eye redness and eye pain, but is only really bothered by the eye pain, each symptom still adds to the overall score by 1.

Another study shows TEPEZZA is proven to help patients who were diagnosed with TED years ago



6 out of 10 patients who took TEPEZZA saw **reduced eye bulging*** vs 2 out of 10 people taking placebo†

Individual results may vary.
*Eye bulging was reduced by at least 2 mm.
†All patients in the 24-week study had TED for 2 to 10 years, experienced eye bulging, and didn't have symptom changes for at least one year prior.

Hear real TEPEZZA patients share their experience

TEPEZZA safety information

As with all medications, TEPEZZA may cause side effects, and each person's experience is different. Talk to your doctor about having a plan in place to monitor possible side effects.

SIDE EFFECT	% of patients‡ who experienced the side effect in clinical studies
Muscle spasms	25%
Nausea	17%
Hair loss	13%
Diarrhea	12%
Feeling tired	12%
High blood sugar	10%
Hearing problems	10%
Taste changes	8%
Headache	8%
Dry skin	8%
Weight decrease	6%
Nail disorder	5%

‡These are the most common side effects as reported with TEPEZZA seen in clinical study of patients recently diagnosed with moderate-to-severe TED. Contraindications: TEPEZZA has no listed contraindications.

Changes in menstruation: In clinical studies, about 23% (5 out of 22) of menstruating women saw changes in menstruation. These changes included missing period(s), spotting, and painful menstruation or cramps.

Hearing impairment is one of the side effects reported with TEPEZZA, and it's a topic many patients want to understand. Scan here to learn more.

IMPORTANT SAFETY INFORMATION (cont'd)

Before receiving TEPEZZA, tell your doctor if you:

- Have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- Are currently being treated for diabetes, have been diagnosed with diabetes, or know your blood sugar is high.
- Are pregnant or plan to become pregnant. TEPEZZA may harm your unborn baby. Tell your doctor if you become pregnant or suspect you are pregnant during treatment with TEPEZZA.

Please see additional Important Safety Information for TEPEZZA throughout and on pages 12-13.

TEPEZZA®
teprotumumab-trbw

Learn how TEPEZZA is administered

TEPEZZA is an intravenous (IV) medicine. This means it is delivered through a needle that is placed in your arm. IV medicines like TEPEZZA are most often given by a registered nurse (RN), nurse practitioner (NP), or another medical professional. The process of delivering an IV medicine is called an infusion.

To make the most of your treatment with TEPEZZA, it's recommended that you get all 8 infusions as prescribed by your doctor

Scan here to learn about getting started with TEPEZZA

TEPEZZA treatment includes 8 infusions, one given every 3 weeks.

Your first and second infusions of TEPEZZA will last about 90 minutes each. Your remaining 6 infusions will take about 60 minutes each but could be longer depending on how you react to the infusion.

This is the dosing schedule as prescribed for a patient who reacts well to the initial 2 infusions:



IMPORTANT SAFETY INFORMATION (cont'd)

Before receiving TEPEZZA, tell your doctor if you:

- Women who are able to become pregnant should use an effective form of birth control (contraception) prior to starting treatment, during treatment and for at least 6 months after the final dose of TEPEZZA.
- Are breastfeeding or plan to breastfeed. It is not known if TEPEZZA passes into your breast milk. Talk to your doctor about the best ways to feed your baby during treatment with TEPEZZA.
- Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, dietary and herbal supplements. Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

"After my last infusion, I woke up without double vision. I raced to the mirror because I felt my eyes working."

— Jeanne T., real TEPEZZA patient

Jeanne has been compensated for her time.

Please see additional Important Safety Information for TEPEZZA throughout and on pages 12-13.

USE and IMPORTANT SAFETY INFORMATION

USE

TEPEZZA is a prescription medicine used to treat Thyroid Eye Disease (TED), no matter if you've had TED for months or years.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TEPEZZA?

Infusion reactions can happen during or within 24 hours after your infusion of TEPEZZA. If you have a reaction while receiving TEPEZZA, your doctor or nurse will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction, your doctor may stop your treatment completely.

Tell your doctor or nurse right away if you have any of these symptoms during or after your treatment with TEPEZZA:

- High blood pressure
- Fast heartbeat
- Redness of the face/Feeling hot
- Difficulty breathing
- Headache
- Muscle pain

TEPEZZA may worsen inflammatory bowel disease (IBD), such as Crohn's disease or ulcerative colitis, even if you don't have a diagnosis of IBD before starting TEPEZZA treatment. Tell your doctor right away if you have worsening IBD symptoms, which may include diarrhea with stomach pain or cramps, blood in your stools, sudden urgency to have a bowel movement, or feeling like you need to have a bowel movement even when there's little or no stool to pass.

TEPEZZA may cause an increase in your blood sugar. Before starting treatment with TEPEZZA, tell your doctor if you are currently being treated for diabetes, know your blood sugar is high, or have been diagnosed with diabetes. It is important for you to take your treatments and follow an appropriate diet for glucose control as prescribed by your doctor.

TEPEZZA may cause severe hearing problems including hearing loss, which in some cases may be permanent. Tell your doctor if you have any signs or symptoms of hearing problems or changes in hearing.

Before receiving TEPEZZA, tell your doctor if you:

- Have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- Are currently being treated for diabetes, have been diagnosed with diabetes, or know your blood sugar is high.
- Are pregnant or plan to become pregnant. TEPEZZA may harm your unborn baby. Tell your doctor if you become pregnant or suspect you are pregnant during treatment with TEPEZZA.
    - Women who are able to become pregnant should use an effective form of birth control (contraception) prior to starting treatment, during treatment and for at least 6 months after the final dose of TEPEZZA.
- Are breastfeeding or plan to breastfeed. It is not known if TEPEZZA passes into your breast milk. Talk to your doctor about the best ways to feed your baby during treatment with TEPEZZA.
- Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, dietary and herbal supplements. Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

What are the possible side effects of TEPEZZA?

The most common side effects of TEPEZZA include muscle cramps or spasms, nausea, hair loss, diarrhea, feeling tired, high blood sugar, hearing problems, taste changes, headache, dry skin, ear discomfort, weight loss, nail problems, and changes in menstruation.

This is not a complete list of all possible side effects. Tell your doctor or treatment team about any side effect you may have.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/safety/medwatch, or call 1-800-FDA-1088.

Please see Full Prescribing Information or visit TEPEZZA.com for more information.

TEPEZZA®
teprotumumab-trbw

TEPEZZA is for adults with Thyroid Eye Disease (TED), no matter if they've had TED for months or years

Have even ONE TED SYMPTOM?
like **BULGING EYES** or **DOUBLE VISION**

And even ONE TED IMPACT?
like **FEELING LESS CONFIDENT** or **TROUBLE DRIVING**

Then speak up and ask your doctor about TEPEZZA

Scan here to learn about getting started with TEPEZZA

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TEPEZZA?

Infusion reactions can happen during or within 24 hours after your infusion of TEPEZZA. If you have a reaction while receiving TEPEZZA, your doctor or nurse will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction, your doctor may stop your treatment completely.

Please see additional Important Safety Information for TEPEZZA throughout and on pages 12-13.

AMGEN

© 2026 Amgen Inc. All rights reserved. USA-632-81767 09/26